

History Tracking Report: 2009 to 2008 Requirements

Accreditation Program: Ambulatory Health Care

Chapter: National Patient Safety Goals

NPSG.01.01.01

2009 Requirement Text:

Use at least two [patient] identifiers when providing care, treatment, or services.

Requirement 1A

2008 Requirement Text:

Use at least two {jc}patient{/1} identifiers when providing care, treatment or services.

NPSG.01.01.01

2009 EP: 2

2009 EP Text:

Two patient identifiers are used when administering medications, blood, or blood components.

Requirement 1A

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. Two {jc}patient{/1} identifiers are used when administering medications or blood products.

NPSG.01.01.01

2009 EP: 3

2009 EP Text:

Two patient identifiers are used when collecting blood samples and other specimens for clinical testing.

Requirement 1A

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. Two {jc}patient{/1} identifiers are used when collecting blood samples and other specimens for clinical testing.

NPSG.01.01.01

2009 EP: 4

2009 EP Text:

Two patient identifiers are used when providing other treatments or procedures.

Requirement 1A

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. Two {jc}patient{/1} identifiers are used when providing other treatments or procedures.

NPSG.01.01.01

2009 EP: 5

2009 EP Text:

The patient's room number or physical location is not used as an identifier. (See also MM.05.01.09, EPs 8 and 11)

Requirement 1A

2008 EP: 4

2008 EP Text:

Revision Code: Retain

4. The {jc}patient's{/9} room number or physical location is not used as an identifier.

NPSG.01.01.01

2009 EP: 6

2009 EP Text:

Containers used for blood and other specimens are labeled in the presence of the patient.

Requirement 1A

2008 EP: 5

2008 EP Text:

Revision Code: Retain

5. Containers used for blood and other specimens are labeled in the presence of the {jc}patient{/1}.

NPSG.02.01.01

2009 Requirement Text:

For verbal or telephone orders or for telephone reporting of critical test results, the individual giving the order or test result verifies the complete order or test result by having the person receiving the information record and "read-back" the complete order or test result.

Requirement 2A

2008 Requirement Text:

For verbal or telephone orders or for telephonic reporting of critical test results, verify the complete order or test result by having the person receiving the information record and "read-back" the complete order or test result.

NPSG.02.01.01

2009 EP: 1

2009 EP Text:

The individual receiving the information writes down the complete order or test result or enters it into a computer.

Requirement 2A

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. The receiver of the information writes down the complete order or test result or enters it into a computer.

NPSG.02.01.01

2009 EP: 2

2009 EP Text:

The individual receiving the information reads back the complete order or test result.

Requirement 2A

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The receiver of the information reads back the order or test result.

NPSG.02.01.01

2009 EP: 3

2009 EP Text:

The individual who gave the order or test result confirms the information that was read back.

Requirement 2A

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. The receiver of the information receives confirmation from the individual who gave the order or test result.

NPSG.02.02.01

2009 Requirement Text:

There is a standardized list of abbreviations, acronyms, symbols, and dose designations that are not to be used throughout the [organization].

Requirement 2B

2008 Requirement Text:

Standardize a list of abbreviations, acronyms, symbols, and dose designations that are not to be used throughout the organization.

NPSG.02.02.01

2009 EP: 1

2009 EP Text:

The organization develops a standardized list of abbreviations, acronyms, symbols, and dose designations that are not to be used throughout the organization.

Requirement 2B

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. The organization develops a standardized a list of abbreviations, acronyms, symbols, and dose designations that are not to be used throughout the organization.

NPSG.02.02.01

2009 EP: 2

2009 EP Text:

The current list of abbreviations, acronyms, symbols, and dose designations not to be used includes the following:

- U,u
- IU
- Q.D., QD, q.d., qd
- Q.O.D., QOD, q.o.d, qod
- Trailing zero (X.0 mg)
- Lack of leading zero (.X mg)
- MS
- MSO4
- MgSO4

Note: A trailing zero may be used only when required to demonstrate the level of precision of the value being reported, such as for laboratory results, imaging studies that report the size of lesions, or catheter/tube sizes. It may not be used in medication orders or other medication-related documentation.

Requirement 2B

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The list of abbreviations not to be used includes the following: U,uIU Q.D., QD, q.d., qd Q.O.D., QOD, q.o.d, qodTrailing zero (X.0 mg)*Lack of leading zero (.X mg)MSMSO4MgSO4* Exception: A “trailing zero” may be used only where required to demonstrate the level of precision of the value being reported, such as for laboratory results, imaging studies that report size of lesions, or catheter/tube sizes. It may not be used in medication orders or other medication-related documentation.

NPSG.02.02.01

2009 EP: 3

2009 EP Text:

The organization implements the “do not use” list of abbreviations, acronyms, symbols, and dose designations and applies it to all orders and all medication-related documentation that is handwritten or entered as free text into a computer.

Requirement 2B

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. The organization implements the “do not use” list and applies this list to all orders and all medication-related documentation when handwritten or entered as free text into a computer.

NPSG.02.02.01

2009 EP: 4

2009 EP Text:

The organization does not include any abbreviations, acronyms, symbols, and dose designations identified as not to be used on preprinted forms.

Requirement 2B

2008 EP: 4

2008 EP Text:

Revision Code: Retain

4. Preprinted forms do not include any abbreviations identified as not to be used.

NPSG.02.03.01

2009 Requirement Text:

The [organization] measures, assesses, and, if needed, takes action to improve the timeliness of reporting, and the timeliness of receipt of critical tests and critical results and values by the responsible licensed caregiver.

Requirement 2C

2008 Requirement Text:

Measure, assess, and if appropriate, take action to improve the timeliness of reporting, and the timeliness of receipt by the responsible licensed caregiver, of critical tests and critical results and values.

NPSG.02.03.01

2009 EP: 1

2009 EP Text:

The organization defines critical tests and critical results and values.

Requirement 2C

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1.The organization defines critical tests and critical results and values.

NPSG.02.03.01

2009 EP: 2

2009 EP Text:

The organization defines the acceptable length of time between the ordering of critical tests and reporting the results of these tests, whether normal or abnormal.

Requirement 2C

2008 EP: 2

2008 EP Text:

Revision Code: Split

2. The organization defines the acceptable length of time between the ordering of critical tests and reporting the critical tests and critical results and values.

NPSG.02.03.01

2009 EP: 3

2009 EP Text:

The organization defines the acceptable length of time for reporting the results of routine tests with critical abnormal values or findings.

Requirement 2C

2008 EP: 2

2008 EP Text:

Revision Code: Split

2. The organization defines the acceptable length of time between the ordering of critical tests and reporting the critical tests and critical results and values.

NPSG.02.03.01

2009 EP: 4

2009 EP Text:

The organization defines the acceptable length of time between the availability of critical tests and critical results and values and receipt by the responsible licensed caregiver.

Requirement 2C

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. The organization defines the acceptable length of time between the availability of critical tests and critical results and values and receipt by the responsible licensed care giver.

NPSG.02.03.01

2009 EP: 5

2009 EP Text:

The organization collects data on the timeliness of reporting critical test results and critical results and values from routine tests.

Requirement 2C

2008 EP: 4

2008 EP Text:

Revision Code: Retain

4. The organization collects data on the timeliness of reporting critical tests and critical results and values.

NPSG.02.03.01

2009 EP: 6

2009 EP Text:

The organization assesses the data on the timeliness of reporting critical test results and critical results and values from routine tests and determines whether a need for improvement exists.

Requirement 2C

2008 EP: 5

2008 EP Text:

Revision Code: Retain

5. The organization assesses the data and determines whether there is a need for improvement.

NPSG.02.03.01

2009 EP: 7

2009 EP Text:

The organization takes appropriate action to improve the timeliness of reporting critical test results and critical results and values from routine tests and measures the effectiveness of those actions.

Requirement 2C

2008 EP: 6

2008 EP Text:

6. The organization takes appropriate action to improve and measure the effectiveness of those actions.

Revision Code: Retain

NPSG.02.05.01

2009 Requirement Text:

The [organization] implements a standardized approach to hand-off communications, including an opportunity to ask and respond to questions.

Requirement 2E

2008 Requirement Text:

Implement a standardized approach to “hand-off” communications, including an opportunity to ask and respond to questions.

NPSG.02.05.01

2009 EP: 1

2009 EP Text:

The organization’s process for effective hand-off communication includes the following: Interactive communications that allows for the opportunity for questioning between the giver and receiver of patient information.

Requirement 2E

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. The organization’s process for effective “hand off” communication includes: Interactive communications allowing for the opportunity for questioning between the giver and receiver of {jc}patient{/1} information.

NPSG.02.05.01

2009 EP: 2

2009 EP Text:

The organization’s process for effective hand-off communication includes the following: Up-to-date information regarding the patient’s condition, care, treatment, medications, services, and any recent or anticipated changes. (See also NPSG.08.01.01, EP 4)

Requirement 2E

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The organization’s process for effective “hand off” communication includes: Up-to-date information regarding the {jc}patient’s{/9} care, treatment and services, condition and any recent or anticipated changes.

NPSG.02.05.01

2009 EP: 3

2009 EP Text:

The organization’s process for effective hand-off communication includes the following: A method to verify the received information, including repeat-back or read-back techniques.

Requirement 2E

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. The organization’s process for effective “hand off” communication includes: A process for verification of the received information, including repeat-back or read-back, as appropriate.

NPSG.02.05.01

2009 EP: 4

2009 EP Text:

The organization’s process for effective hand-off communication includes the following: An opportunity for the receiver of the hand-off information to review relevant patient historical data, which may include previous care, treatment, or services.

Requirement 2E

2008 EP: 4

2008 EP Text:

Revision Code: Retain

4. The organization’s process for effective “hand off” communication includes: An opportunity for the receiver of the hand off information to review relevant {jc}patient{/1} historical data, which may include previous care, treatment and services.

NPSG.02.05.01

2009 EP: 5

2009 EP Text:

Interruptions during hand-offs are limited to minimize the possibility that information fails to be conveyed or is forgotten.

Requirement 2E

2008 EP: 5

2008 EP Text:

Revision Code: Retain

5. Interruptions during hand offs are limited to minimize the possibility that information would fail to be conveyed or would be forgotten.

NPSG.03.03.01

2009 Requirement Text:

The [organization] identifies and, at a minimum, annually reviews a list of look-alike/sound-alike medications used by the [organization] and takes action to prevent errors involving the interchange of these medications.

Requirement 3C

2008 Requirement Text:

Identify and, at a minimum, annually review a list of look-alike/sound-alike drugs used by the organization, and take action to prevent errors involving the interchange of these drugs.

NPSG.03.03.01

2009 EP: 1

2009 EP Text:

The organization identifies a list of look-alike/sound-alike medications used by the organization. The list includes a minimum of 10 look-alike/sound-alike medication combinations selected from the tables of look-alike/sound-alike medications posted on The Joint Commission Web site at <http://www.jointcommission.org>.

Requirement 3C

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. Identify a list of look-alike/sound-alike (LASA) drugs used by the organization (the list must include a minimum of 10 look-alike/sound-alike drug combinations selected from the tables of LASA drugs posted on the Joint Commission website).

NPSG.03.03.01

2009 EP: 2

2009 EP Text:

The organization reviews the list of look-alike/sound-alike medications at least annually.

Requirement 3C

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. Review the list of look-alike/sound-alike drugs used by the organization at least annually.

NPSG.03.03.01

2009 EP: 3

2009 EP Text:

The organization takes action to prevent errors involving the interchange of the medications on the list of look-alike/sound-alike medication list.

Requirement 3C

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. The organization takes action to prevent errors involving the interchange of these drugs.

NPSG.03.04.01**2009 Requirement Text:**

Label all medications, medication containers (for example, syringes, medicine cups, basins), or other solutions on and off the sterile field.

Requirement 3D**2008 Requirement Text:**

Label all medications, medication containers (for example, syringes, medicine cups, basins), or other solutions on and off the sterile field.

NPSG.03.04.01**2009 EP: 1****2009 EP Text:**

Medications and solutions both on and off the sterile field are labeled even if there is only one medication being used.

Requirement 3D**2008 EP: 1****2008 EP Text:****Revision Code:** Retain

1. Medications and solutions both on and off the sterile field are labeled even if there is only one medication being used.

NPSG.03.04.01**2009 EP: 2****2009 EP Text:**

Labeling occurs when any medication or solution is transferred from the original packaging to another container.

Requirement 3D**2008 EP: 2****2008 EP Text:****Revision Code:** Retain

2. Labeling occurs when any medication or solution is transferred from the original packaging to another container.

NPSG.03.04.01**2009 EP: 3****2009 EP Text:**

Medication or solution labels include the medication name, strength, amount (if not apparent from the container), expiration date when not used within 24 hours, and expiration time when expiration occurs in less than 24 hours.

Requirement 3D**2008 EP: 3****2008 EP Text:****Revision Code:** Retain

3. Labels include the drug name, strength, amount (if not apparent from the container), expiration date when not used within 24 hours, and expiration time when expiration occurs in less than 24 hours.

NPSG.03.04.01**2009 EP: 4****2009 EP Text:**

All medication or solution labels are verified both verbally and visually by two qualified individuals whenever the person preparing the medication or solution is not the person who will be administering it.

Requirement 3D**2008 EP: 4****2008 EP Text:****Revision Code:** Retain

4. All labels are verified both verbally and visually by two qualified individuals when the person preparing the medication is not the person administering the medication.

NPSG.03.04.01**2009 EP: 5****2009 EP Text:**

No more than one medication or solution is labeled at one time.

Requirement 3D**2008 EP: 5****2008 EP Text:****Revision Code:** Retain

5. No more than one medication or solution is labeled at one time.

NPSG.03.04.01**2009 EP: 6****2009 EP Text:**

Any medications or solutions found unlabeled are immediately discarded.

Requirement 3D**2008 EP: 6****2008 EP Text:****Revision Code:** Retain

6. Any medications or solutions found unlabeled are immediately discarded.

NPSG.03.04.01	2009 EP: 7	Requirement 3D	2008 EP: 7
2009 EP Text:		2008 EP Text:	Revision Code: Retain
All original containers from medications or solutions remain available for reference in the perioperative or procedural area until the conclusion of the procedure.		7.All original containers from medications or solutions remain available for reference in theperioperative/procedural area until the conclusion of the procedure.	
NPSG.03.04.01	2009 EP: 8	Requirement 3D	2008 EP: 8
2009 EP Text:		2008 EP Text:	Revision Code: Retain
All labeled containers on the sterile field are discarded at the conclusion of the procedure.		8. All labeled containers on the sterile field are discarded at the conclusion of the procedure.	
NPSG.03.04.01	2009 EP: 9	Requirement 3D	2008 EP: 9
2009 EP Text:		2008 EP Text:	Revision Code: Retain
At shift change or break relief, all medications and solutions both on and off the sterile field and their labels are reviewed by entering and exiting personnel.		9. At shift change or break relief, all medications and solutions both on and off the sterile field and their labels are reviewed by entering and exiting personnel	

NPSG.03.05.01

2009 Requirement Text:

Reduce the likelihood of [patient] harm associated with the use of anticoagulation therapy.

Note: This requirement applies only to [organization]s that provide anticoagulation therapy and/or long-term anticoagulation prophylaxis (for example, atrial fibrillation) where the clinical expectation is that the [patient]'s laboratory values for coagulation will remain outside normal values. This requirement does not apply to routine situations where short-term prophylactic anticoagulation is used for venous thrombo-embolism prevention (for example, related to procedures or hospitalization) and the clinical expectation is that the [patient]'s laboratory values for coagulation will remain within, or close to, normal values.

Requirement 3E

2008 Requirement Text:

Reduce the likelihood of patient harm associated with the use of anticoagulation therapy. Note: This requirement applies only to organizations that provide anticoagulation therapy.

NPSG.03.05.01

2009 EP: 1

2009 EP Text:

The organization implements a defined anticoagulation management program to individualize the care provided to each patient receiving anticoagulant therapy.

Requirement 3E

2008 EP: 5

2008 EP Text:

Revision Code: Retain

5. The organization implements a defined anticoagulant management program to individualize the care provided to each patient receiving anticoagulant therapy.

NPSG.03.05.01

2009 EP: 2

2009 EP Text:

To reduce compounding and labeling errors, the organization uses only oral unit dose products, pre-filled syringes, or pre-mixed infusion bags when these types of products are available.

Note: For pediatric patients, pre-loaded syringe products should only be used if specifically designed for children.

Requirement 3E

2008 EP: 6

2008 EP Text:

Revision Code: Retain

6. To reduce compounding and labeling errors, the organization uses ONLY oral unit dose products and pre-mixed infusions, when these products are available.

NPSG.03.05.01

2009 EP: 3

2009 EP Text:

The organization uses approved protocols for the initiation and maintenance of anticoagulation therapy appropriate to the medication used, to the condition being treated, and to the potential for medication interactions.

Requirement 3E

2008 EP: 8

2008 EP Text:

Revision Code: Retain

8. The organization uses approved protocols for the initiation and maintenance of anticoagulation therapy appropriate to the medication used, to the condition being treated, and to the potential for drug interactions.

NPSG.03.05.01

2009 EP: 4

2009 EP Text:

For patients starting on warfarin, a baseline International Normalized Ratio (INR) is available, and for all patients receiving warfarin therapy, a current INR is available and is used to monitor and adjust therapy.

Requirement 3E

2008 EP: 9

2008 EP Text:

Revision Code: Retain

9. For patients being started on warfarin, a baseline International Normalized Ratio (INR) is available, and for all patients receiving warfarin therapy, a current INR is available and is used to monitor and adjust therapy.

NPSG.03.05.01	2009 EP: 5	Requirement 3E	2008 EP: 10
2009 EP Text:		2008 EP Text:	Revision Code: Retain
When dietary services are provided by the organization, the service is notified of all patients receiving warfarin and responds according to its established food/medication interaction program.		10. When dietary services are provided by the organization, the service is notified of all patients receiving warfarin and responds according to its established food/drug interaction program.	
NPSG.03.05.01	2009 EP: 6	Requirement 3E	2008 EP: 11
2009 EP Text:		2008 EP Text:	Revision Code: Retain
When heparin is administered intravenously and continuously, the organization uses programmable infusion pumps in order to provide consistent and accurate dosing.		11. When heparin is administered intravenously and continuously, the organization uses programmable infusion pumps.	
NPSG.03.05.01	2009 EP: 7	Requirement 3E	2008 EP: 12
2009 EP Text:		2008 EP Text:	Revision Code: Retain
The organization has a written policy that addresses baseline and ongoing laboratory tests that are required for heparin and low molecular weight heparin therapies.		12. The organization has a policy that addresses baseline and ongoing laboratories tests that are required for heparin and low molecular weight heparin therapies.	
NPSG.03.05.01	2009 EP: 8	Requirement 3E	2008 EP: 13
2009 EP Text:		2008 EP Text:	Revision Code: Retain
The organization provides education regarding anticoagulation therapy to staff, patients, and families. Note: Patient/family education includes the importance of follow-up monitoring, compliance issues, dietary restrictions, and potential for adverse drug reactions and interactions.		13. The organization provides education regarding anticoagulation therapy to staff, patients, and families.	
NPSG.03.05.01	2009 EP: 9	Requirement 3E	2008 EP: 15
2009 EP Text:		2008 EP Text:	Revision Code: Retain
The organization evaluates its anticoagulation safety practices, takes appropriate action to improve its practices, and measures the effectiveness of those actions on a regular basis.		15. The organization evaluates anticoagulation safety practices (see MM.8.10).	

NPSG.07.01.01

2009 Requirement Text:

Comply with current World Health Organization (WHO) hand hygiene guidelines or Centers for Disease Control and Prevention (CDC) hand hygiene guidelines.

Requirement 7A

2008 Requirement Text:

Comply with current World Health Organization (WHO) Hand Hygiene Guidelines or Centers for Disease Control and Prevention (CDC) hand hygiene guidelines.

NPSG.07.01.01

2009 EP: 1

2009 EP Text:

The organization complies with current World Health Organization (WHO) or Centers for Disease Control and Prevention (CDC) hand hygiene guidelines.
 Note: Organizations are required to comply with 1A, 1B, and 1C of the WHO or CDC guidelines.

Requirement 7A

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. Comply with current WHO Hand Hygiene Guidelines or Centers for Disease Control and Prevention (CDC) hand hygiene guidelines* *Organizations are required to comply with all 1A, 1B, 1C CDC or WHO guidelines.

NPSG.07.02.01

2009 Requirement Text:

Manage as sentinel events all identified cases of unanticipated death or major permanent loss of function related to a health care associated infection.

Requirement 7B

2008 Requirement Text:

Manage as sentinel events all identified cases of unanticipated death or major permanent loss of function associated with a health care-associated infection.

NPSG.07.02.01

2009 EP: 1

2009 EP Text:

The organization manages all identified cases of unanticipated death or major permanent loss of function associated with a health care associated infection as sentinel events (that is, the organization conducts a root cause analysis).

Requirement 7B

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. The organization manages all identified cases of unanticipated death or major permanent loss of function associated with a health care-associated infection as sentinel events (that is, conducts a root cause analysis).

NPSG.07.02.01

2009 EP: 2

2009 EP Text:

The root cause analysis addresses the management of the patient before and after the identification of infection.

Requirement 7B

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The root cause analysis addresses the management of the {jc}patient{/1} before and after the identification of infection.

NPSG.08.01.01

2009 Requirement Text:

A process exists for comparing the [patient]'s current medications with those ordered for the [patient] while under the care of the [organization].

Requirement 8A

2008 Requirement Text:

There is a process for comparing the {jc}patient's{/9} current medications with those ordered for the {jc}patient{/1} while under the care of the organization.

NPSG.08.01.01

2009 EP: 1

2009 EP Text:

At the time the patient enters the organization or is admitted, a complete list of the medications the patient is taking at home (including dose, route, and frequency) is created and documented. The patient, and family as needed, are involved in creating this list.

Requirement 8A

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. The organization, with the {jc}patient's{/9} involvement, creates a complete list of the {jc}patient's{/9} current medications at admission/entry.

NPSG.08.01.01

2009 EP: 2

2009 EP Text:

The medications ordered for the patient while under the care of the organization are compared to those on the list created at the time of entry to the organization or admission.

Requirement 8A

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The medications ordered for, administered to, or dispensed to the {jc}patient{/1} while under the care of the organization are compared to those on the list and any discrepancies (e.g., omissions, duplications, potential interactions) are resolved.

NPSG.08.02.01

2009 Requirement Text:

When a [patient] is referred to or transferred from one [organization] to another, the complete and reconciled list of medications is communicated to the next provider of service and the communication is documented. Alternatively, when a [patient] leaves the [organization]’s care directly to his or her home, the complete and reconciled list of medications is provided to the [patient]’s known primary care provider, or the original referring provider, or a known next provider of service.

Note: When the next provider of service is unknown or when no known formal relationship is planned with a next provider, giving the [patient], and family as needed, the list of reconciled medications is sufficient.

Requirement 8B

2008 Requirement Text:

A complete list of the {jc}patient’s{/9} medications is communicated to the next provider of service when a {jc}patient{/1} is referred or transferred to another setting, service, practitioner or level of care within or outside the organization. The complete list of medications is also provided to the patient on discharge from the facility.

NPSG.08.02.01

2009 EP: 1

2009 EP Text:

The patient’s most current reconciled medication list is communicated to the next provider of service, either within or outside the organization. The communication between providers is documented.

Requirement 8B

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. The {jc}patient{/1}’s accurate medication reconciliation list (complete with medications prescribed by the first provider of service) is communicated to the next provider of service, whether it be within or outside the organization

NPSG.11.01.01

2009 Requirement Text:

The [organization] educates staff, including licensed independent practitioners who are involved with surgical procedures and anesthesia providers, on how to control heat sources and how to manage fuels while maintaining enough time for [patient] preparation, and establishes guidelines to minimize oxygen concentration under drapes.

Requirement 11A

2008 Requirement Text:

Educate staff, including operating licensed independent practitioners and anesthesia providers, on how to control heat sources and manage fuels with enough time for {jc}patient{/1} preparation, and establish guidelines to minimize oxygen concentration under drapes.

NPSG.11.01.01

2009 EP: 1

2009 EP Text:

Organizations assess the risk for surgical fires based on equipment and procedures used.

Requirement 11A

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. Organizations assess the risk for surgical fires based on equipment and procedures used.

NPSG.11.01.01

2009 EP: 2

2009 EP Text:

The organization establishes guidelines to minimize oxygen concentrations under drapes.

Requirement 11A

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The organization establishes guidelines to minimize oxygen concentrations under drapes.

NPSG.11.01.01

2009 EP: 3

2009 EP Text:

Organizations that identify themselves at risk for surgical fires provide staff training on methods to minimize oxygen concentration under drapes.

Requirement 11A

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. Organizations that identify themselves at risk provide staff training on methods to minimize oxygen concentration under drapes

NPSG.11.01.01

2009 EP: 4

2009 EP Text:

Organizations that identify themselves at risk for surgical fires provide staff training on methods to avoid the use of flammable solutions and materials.

Requirement 11A

2008 EP: 4

2008 EP Text:

Revision Code: Retain

4. Organizations that identify themselves at risk provide staff training on methods to avoid the use of flammable solutions and materials.

NPSG.11.01.01

2009 EP: 5

2009 EP Text:

Organizations that identify themselves at risk for surgical fires provide staff training on actions to take in the event of a surgical fire.

Requirement 11A

2008 EP: 5

2008 EP Text:

Revision Code: Retain

5. Organizations that identify themselves at risk provide staff training on actions to take in the event of a surgical fire.

NPSG.13.01.01

2009 Requirement Text:

Identify the ways in which the [patient] and his or her family can report concerns about safety and encourage them to do so.

Requirement 13A

2008 Requirement Text:

Define and communicate the means for {jc}patients{/6} and their families to report concerns about safety and encourage them to do so.

NPSG.13.01.01

2009 EP: 1

2009 EP Text:

The patient and family are educated on available reporting methods for concerns related to care, treatment, services and patient safety issues.

Requirement 13A

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. {jc}Patients{/6} and families are educated on methods available to report concerns related to care, treatment, services and {jc}patient{/1} safety issues.

NPSG.13.01.01

2009 EP: 4

2009 EP Text:

The organization encourages patients and their families to report concerns about safety.

Requirement 13A

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The organization encourages {jc}patient{/1}s and their families to report concerns about safety.

UP.01.01.01

2009 Requirement Text:

Conduct a pre-procedure verification process.

UP Requirement 1A

2008 Requirement Text:

Conduct a pre-operative verification process as described in the Universal Protocol

UP.01.01.01

2009 EP: 1

2009 EP Text:

Verification of the correct person, correct site, and correct procedure occurs at the following times:

- At the time the procedure is scheduled.
- At the time of preadmission testing and assessment.
- At the time of admission or entry into the facility for a procedure, whether elective or emergent.
- Before the patient leaves the pre-procedure area or enters the procedure room.
- Anytime the responsibility for care of the patient is transferred to another member of the procedural care team, (including the anesthesia providers) at the time of, and during, the procedure.
- With the patient involved, awake and aware, if possible.

UP Requirement 1A

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. Verification of the correct person, procedure, and site should occur during the following (as applicable): At the time the surgery/procedure is scheduled. At the time of admission or entry into the facility. Anytime the responsibility for care of the patient is transferred to another caregiver. With the patient involved, awake and aware, if possible. Before the patient leaves the preoperative area or enters the procedure/surgical room.

UP.01.01.01

2009 EP: 2

2009 EP Text:

When the patient is in the pre-procedure area, immediately prior to moving the patient to the procedure room, a checklist (for example, paper, electronic, or other medium such as a wall-mounted white-board) is used to review and verify that the following items are available and accurately matched to the patient:

- Relevant documentation (for example, history and physical, nursing assessment, and pre-anesthesia assessment).
- Accurately completed, and signed, procedure consent form.
- Correct diagnostic and radiology test results (for example, radiology images and scans, or pathology and biopsy reports) that are properly labeled.
- Any required blood products, implants, devices and/or special equipment for the procedure.

UP Requirement 1A

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The following is reviewed prior to the start of the procedure: Relevant documentation (e.g. H&P, consent). Relevant images, properly labeled and displayed. Any required implants and special equipment.

UP.01.02.01

2009 Requirement Text:

Mark the procedure site.

UP Requirement 1B

2008 Requirement Text:

Mark the operative site as described in the Universal Protocol

UP.01.02.01

2009 EP: 1

2009 EP Text:

For all procedures involving incision or percutaneous puncture or insertion, the intended procedure site is marked. The marking takes into consideration laterality, the surface (flexor, extensor), the level (spine), or specific digit or lesion to be treated.

Note: For procedures that involve laterality of organs but the incision(s) or approaches may be from the mid-line or from a natural orifice, the site is still marked and the laterality noted.

UP Requirement 1B

2008 EP: 5

2008 EP Text:

Revision Code: Retain

5. At a minimum, mark all cases involving laterality, multiple structures (fingers, toes, lesions), or multiple levels (spine). (Note: In addition to pre-operative skin marking of the general spinal region, special intraoperative radiographic techniques are used for marking the exact vertebral level).

UP.01.02.01

2009 EP: 2

2009 EP Text:

The procedure site is initially marked before the patient is moved to the location where the procedure will be performed and takes place with the patient involved, awake and aware, if possible.

UP Requirement 1B

2008 EP: 7

2008 EP Text:

Revision Code: Retain

7. Marking must take place with the patient involved, awake and aware, if possible.

UP.01.02.01

2009 EP: 3

2009 EP Text:

The procedure site is marked by a licensed independent practitioner or other provider who is privileged or permitted by the organization to perform the intended surgical or non-surgical invasive procedure. This individual will be involved directly in the procedure and will be present at the time the procedure is performed.

Note: Final confirmation and verification of the site mark takes place during the time-out.

UP Requirement 1B

2008 EP: 6

2008 EP Text:

Revision Code: Retain

6. The person performing the procedure should do the site marking.

UP.01.02.01

2009 EP: 4

2009 EP Text:

The method of marking the site and the type of mark is unambiguous and is used consistently throughout the organization.

UP Requirement 1B

2008 EP: 2

2008 EP Text:

Revision Code: Consolidate

2. The mark must be unambiguous. (Note: for example, use initials or "YES" or a line representing the proposed incision; consider that "X" may be ambiguous.)

<p>UP.01.02.01</p> <p>2009 EP Text:</p> <p>The method of marking the site and the type of mark is unambiguous and is used consistently throughout the organization.</p>	<p>2009 EP: 4</p>	<p>UP Requirement 1B</p> <p>2008 EP Text:</p> <p>4. The method of marking and type of mark should be consistent throughout the organization.</p>	<p>2008 EP: 4</p> <p>Revision Code: Consolidate</p>
<p>UP.01.02.01</p> <p>2009 EP Text:</p> <p>The site marking has the following characteristics:</p> <ul style="list-style-type: none"> - It is made at or near the procedure site or the incision site. Other non-procedure site(s) are not marked unless necessary for some other aspect of care. - It includes, preferably, the surgeon's or proceduralist's initials, with or without a line representing the proposed incision. - It is made using a marker that is sufficiently permanent to remain visible after completion of the skin prep and sterile draping. Adhesive site markers are not to be used as the sole means of marking the site. - It is positioned to be visible after the patient has his or her skin prepped, is in his or her final position, and sterile draping is completed. 	<p>2009 EP: 5</p>	<p>UP Requirement 1B</p> <p>2008 EP Text:</p> <p>3. The mark must be positioned to be visible after the patient is prepped and draped.</p>	<p>2008 EP: 3</p> <p>Revision Code: Consolidate</p>
<p>UP.01.02.01</p> <p>2009 EP Text:</p> <p>The site marking has the following characteristics:</p> <ul style="list-style-type: none"> - It is made at or near the procedure site or the incision site. Other non-procedure site(s) are not marked unless necessary for some other aspect of care. - It includes, preferably, the surgeon's or proceduralist's initials, with or without a line representing the proposed incision. - It is made using a marker that is sufficiently permanent to remain visible after completion of the skin prep and sterile draping. Adhesive site markers are not to be used as the sole means of marking the site. - It is positioned to be visible after the patient has his or her skin prepped, is in his or her final position, and sterile draping is completed. 	<p>2009 EP: 5</p>	<p>UP Requirement 1B</p> <p>2008 EP Text:</p> <p>1. Make the mark at or near the incision site; do not mark any non-operative site(s) unless necessary for some other aspect of care.</p>	<p>2008 EP: 1</p> <p>Revision Code: Consolidate</p>

UP.01.03.01

2009 Requirement Text:

A time-out is performed immediately prior to starting procedures.

UP Requirement 1C

2008 Requirement Text:

Conduct a “time out” immediately before starting the procedure as described in the Universal Protocol

UP.01.03.01

2009 EP: 1

2009 EP Text:

The time-out is conducted prior to starting the procedure and, ideally, prior to the introduction of the anesthesia process (including general/regional anesthesia, local anesthesia, and spinal anesthesia), unless contraindicated.

UP Requirement 1C

2008 EP: 1

2008 EP Text:

Revision Code: Retain

1. The final verification process must be conducted in the location where the procedure will be done, just before starting the procedure.

UP.01.03.01

2009 EP: 5

2009 EP Text:

The time-out addresses the following:

- Correct patient identity.
- Confirmation that the correct side and site are marked.
- An accurate procedure consent form.
- Agreement on the procedure to be done.
- Correct patient position.
- Relevant images and results are properly labeled and appropriately displayed.
- The need to administer antibiotics or fluids for irrigation purposes. (See also NPSG.07.05.01, EP 7)
- Safety precautions based on patient history or medication use.

UP Requirement 1C

2008 EP: 2

2008 EP Text:

Revision Code: Retain

2. The process must involve the entire operative team, use active communication, and must, at least, include: Correct patient identity. Correct side and site Agreement on the procedure to be done. Correct patient position. Availability of correct implants and any special equipment or special requirements.

UP.01.03.01

2009 EP: 6

2009 EP Text:

The completed components of the Universal Protocol and time-out are clearly documented.

UP Requirement 1C

2008 EP: 3

2008 EP Text:

Revision Code: Retain

3. The process is briefly documented, such as in a checklist (Note: the organization should determine the type and amount of documentation.)