

2004 Home Care National Patient Safety Goals

The Joint Commission is "tailoring" the NPSGs to each of its non-hospital programs. This is the first phase of a process to make the NPSGs more relevant to the other programs. The second phase will involve identifying new program-specific evidence- or experience-based goals and requirements. Phase II is planned for implementation beginning in January 2005.

1) Improve the accuracy of patient identification.

- a) Use at least two patient identifiers whenever taking blood samples or administering medications or blood products.
- b) Prior to the start of any surgical or invasive procedure, conduct a final verification process, such as a "time out," to confirm the correct patient, procedure and site, using active—not passive—communication techniques.

2) Improve the effectiveness of communication among caregivers.

- a) Implement a process for taking verbal or telephone orders or critical test results that require a verification "read-back" of the complete order or test result by the person receiving the order or test result.
- b) *Standardize the abbreviations, acronyms and symbols used throughout the organization, including a list of abbreviations, acronyms and symbols not to use.* [Scored at Standard IM.3.10, EP #2]

3) Improve the safety of using high-alert medications.

- a) *Remove concentrated electrolytes (including, but not limited to, potassium chloride, potassium phosphate, sodium chloride >0.9%) from patient care areas.* [Scored at Standard MM.2.20, EP #9]
- b) *Standardize and limit the number of drug concentrations used by the organization.* [Scored at Standard MM.2.20, EP #8]

4) Not applicable

5) Improve the safety of using infusion pumps.

- a) Ensure free-flow protection on all general-use and PCA (patient controlled analgesia) intravenous infusion pumps used by the organization.

6) Improve the effectiveness of clinical alarm systems.

- a) Implement regular preventive maintenance and testing of alarm systems in accordance with manufacturer's guidelines.
- b) Assure that alarms are activated with appropriate settings and are sufficiently audible with respect to distances and competing noise within the patient care area.

7) Reduce the risk of health care-acquired infections

- a) Comply with current U.S. Centers for Disease Control and Prevention (CDC) hand hygiene guidelines.
- b) Manage as sentinel events all identified cases of unanticipated death or major permanent loss of function associated with health-care acquired infection.