

# SpeakUp<sup>\*</sup>

# Speak Up<sup>™</sup> To Prevent Falls



# Take care of your health

- Exercise regularly. It builds strength and some exercises can improve your balance.
- Prevent dehydration. It can affect your balance.
- Have your vision checked.
- If your medicine leaves you drowsy, dizzy, weak or confused, tell your doctor. Ask how to reduce these side effects or if you can take another medicine.



#### Take extra precautions

- Turn on the lights when you enter a room. Do not walk in the dark.
- Make sure your pathway is clear.
- Use the handrails on staircases.
- Sit in chairs that do not move and have arm rests to help you sit down and stand up.
- Wear shoes that have firm, flat, non-slip soles.
- Do not wear shoes that do not have backs on them.
- Replace the rubber tips on canes and walkers when they become worn.

/	5/	$\searrow$	
(		<u>}</u>	).
	Ľ		/
		/	

### Make small changes to your home

- Install timers, "clap-on" or motion sensors on your lights.
- Declutter regular pathways, such as to the bathroom and in poorly lit areas.
- Use night lights in your bedroom, bathroom and the hallway.
- Remove rugs that can slip. Rubber mats are a good replacement.
- Put frequently used items in easy-to-reach places that do not require using a step stool.
- Make sure it is easy to get in and out of your bed.
- Apply non-slip treads on stairs.
- Apply decals or waterproof, non-slip mats in the bathtub and shower.
- Install grab bars near the toilet and the bathtub or shower.
- A home care agency, personal care and support agency, or community program may be able to help make changes to your home if you live alone and need help.



## Ask for help in the hospital or nursing home

- Always use your call button to ask for help getting out of bed. It's for your safety. You may be weaker than you realize.
- Pay attention to what your doctors or nurses tell you about your risk of falling. A fall can mean a longer hospital stay. Also, injuries from a fall can affect your health for the rest of your life.
- Tell your doctor or nurse if your medicine makes you sleepy, light-headed, dizzy, sluggish, unbalanced or confused.
- Do not feel embarrassed asking for help going to the toilet. You will need extra help until you get stronger.
- Wear non-slip socks or footwear.
- Lower the height of the bed and the side rails.

#### The goal of Speak Up™ is to help patients and their advocates become active in their care.

Speak  $Up^{TM}$  materials are intended for the public and have been put into a simplified (i.e., easy-to-read) format to reach a wider audience. They are not meant to be comprehensive statements of standards interpretation or other accreditation requirements, nor are they intended to represent evidence-based clinical practices or clinical practice guidelines. Thus, care should be exercised in using the content of Speak  $Up^{TM}$  materials. Speak  $Up^{TM}$  materials are available to all health care organizations; their use does not indicate that an organization is accredited by The Joint Commission.